



New Zealand standardised oral formulation batch sheet

Tramadol suspension 10 mg/mL (100 mL)

Patient's name		Storage condition	In a refrigerator	At room temperature
NHI		Temperature	2-8°C	≤ 25°C
Date compounded		Shelf life	30 days	30 days
Batch number		Recommended storage	<input checked="" type="checkbox"/>	May be stored at room temperature if this is more practical and temperatures greater than 25°C are avoided

Ingredients required and formula:

Ingredient	Supplier	Batch number	Expiry date	Standard formula	Quantity dispensed	Dispensed by	Checked by
Tramadol 50 mg capsules				20 capsules			
Ora-Blend SF [®]	Perrigo			to 100 mL			

OR

Ora-Plus [®]	Perrigo			50 mL			
Ora-Sweet SF [®]	Perrigo			50 mL			

Ora-Blend[®] or Ora-Sweet[®] can be substituted? No

Calculations performed by

Calculations checked by (pharmacist)

Area cleared for processing by

Instructions to compound suspension:

Note: this formula uses **Ora-Blend SF[®]** or **Ora-Sweet SF[®]**

1. **If Ora-Blend SF[®] unavailable:** Pre-mix the Ora-Plus[®] and Ora-Sweet SF[®] to form the diluent.
2. Empty the contents of the capsules in a mortar.
3. Add diluent (Ora-Blend SF[®] or prepared diluent) to form a smooth paste.
4. Gradually add diluent and transfer to the final measuring flask, rinsing the mortar well. Mix well.
5. Make up to final volume and mix well.
6. Label and package product appropriately.

Labels

Attach product label and auxiliary labels:

Label checked by



New Zealand standardised oral formulation batch sheet

Tramadol suspension 10 mg/mL (100 mL)

Final yield	_____ mL	Checked by	
Final appearance of product	White suspension.		
Container	Amber plastic mission bottle/ amber glass.		
Compounded by	<i>Name:</i>	<i>Signature:</i>	<i>Date:</i>
Final check and product release	<i>Name:</i>	<i>Signature:</i>	<i>Date:</i>
Special instructions: 1. Shake well before use. 2. Store in a refrigerator.			
References: 1. Polonini HC, Silva SL, Cunha CN, et al. <i>Pharmazie</i> , 2016; 71 (4): 185-191. 2. Wagner DS, Johnson CE, Cichon-Hensley BK, et al. <i>Am J Health-Syst Pharm</i> , 2003; 60 : 1268-1270.			
Prepared by	New Zealand Compounding Working Group		
Approved by	Medication Safety Expert Advisory Group of Health Quality and Safety Commission NZ		
Hosted by	Pharmaceutical Society of New Zealand Standard Formulations in the Practice Support section of the PSNZ website		
Version number	3.0	Version approval date	17/07/19
Document review due	01/08/21		

Disclaimer: this batch sheet has been designed to provide guidance and standardised formulations for New Zealand pharmacists. Please read and familiarise yourself with the general guidance for compounding oral liquids (available on the PSNZ web site) before use. Information was considered accurate at the time of publication.